

Mini-review on baryogenesis at the TeV scale and possible connections with dark matter

J. Racker

*Departamento de Física Teórica, Instituto de Física corpuscular (IFIC), Universidad de Valencia-CSIC
Edificio de Institutos de Paterna, Apt. 22085, 46071 Valencia, Spain*

Abstract

This is a very short review on different mechanisms for baryogenesis from particle decays or annihilations at low temperature ($T \lesssim 10$ TeV) and their implementation in models that relate the origin of the baryon asymmetry and dark matter.

Keywords: Baryogenesis, Dark matter, Neutrino masses, Beyond Standard Model

Many scenarios for explaining the origin of the Baryon Asymmetry of the Universe (BAU), notably “standard” leptogenesis, must take place at high temperatures ($T \gtrsim 10^8$ GeV). However there are several motivations to explore models for baryogenesis at lower scales, like accessibility to experimental exploration, compatibility with some well motivated supergravity models that require reheating temperatures below $10^5 - 10^7$ GeV not to overproduce gravitinos [1, 2], and avoidance of hierarchy problems [3]. Moreover, it could also happen that baryogenesis above a few TeVs becomes severely disfavored, demanding a low-scale explanation for the BAU, e.g. if some lepton number (L) violating processes are observed at the LHC [4, 5]. Here we review the main problems to achieve baryogenesis from heavy particle decays or annihilations at low temperatures, mechanisms to reach the TeV scale, and some examples of their use in models connecting the origin of the BAU and Dark Matter (DM).

A crucial quantity for baryogenesis is the amount of CP violation per each decay or annihilation, to be denoted by ϵ . E.g. in type I unflavoured leptogenesis, $\epsilon \equiv \frac{\gamma(N \rightarrow \ell H) - \gamma(N \rightarrow \bar{\ell} \bar{H})}{\gamma(N \rightarrow \ell H) + \gamma(N \rightarrow \bar{\ell} \bar{H})}$, where N is a sterile neutrino, H the Standard Model (SM) Higgs, ℓ a SM lepton doublet, and the γ 's denote number of processes per unit time and volume. To simplify this discussion we take ϵ independent of the temperature (which is often a very good approximation). In this case the baryon density asymmetry, n_B , conveniently normalized to the entropy density, s , is directly proportional to ϵ and can be expressed as $Y_B \equiv \frac{n_B}{s} = \kappa \epsilon \eta$. The “efficiency” η depends

on the kinetics of baryogenesis, while the numerical coefficient κ is defined in each model under the condition that the maximum efficiency be 1. There are two main problems to have baryogenesis at low scales:

- (a) In some leptogenesis models the CP asymmetry ϵ is proportional to the masses of both, the light -mostly active- neutrinos and the heavy ones [6]. Given the tiny value of the light neutrino masses, the mass M of the heavy neutrinos -which sets the scale for baryogenesis- must be very high to have a large enough ϵ . E.g. in type I leptogenesis with hierarchical heavy neutrinos, $M \gtrsim 10^8 - 10^9$ GeV [6, 7]. This fact seems to be well known.
- (b) The second problem is intrinsic to all scenarios of baryogenesis from particle decays or annihilations, but curiously enough it is often either completely ignored or not treated properly. CP violation requires a CP odd and a CP even phase. The former arises from complex couplings, the later from the absorptive part of one loop contributions. In turn the absorptive part is proportional to the amplitude of processes which violate baryon number (B) -or L - [8], and whose strength, therefore, cannot be very small for the CP asymmetry to be large enough. If baryogenesis occurs at low temperatures, when the expansion rate of the universe is mild, these processes are typically very fast and washout the asymmetry.

To illustrate problem (b) consider leptogenesis in a radiative seesaw model -to be safe from problem (a)-, namely the Inert Doublet Model (IDM) with singlet

neutrinos: The SM is extended with a Higgs doublet, H_2 , and at least two heavy sterile neutrinos, N_i ($i = 1, 2$). These extra fields are the only ones odd under a new Z_2 discrete symmetry, which is assumed to be exactly conserved, so that H_2 does not acquire a vacuum expectation value [9]. Choosing the mass of N_1 larger than the mass of H_2 , $M_1 > M_{H_2}$, the CP asymmetry in N_1 decays arises from the interference of tree level and one loop diagrams, as depicted in figure 1. The CP asymmetry ϵ is proportional to the amplitude of the processes at the right of the cut, $H_2 \ell \leftrightarrow \bar{H}_2 \bar{\ell}$ ¹, which may washout the lepton asymmetry depending on how fast they are compared to the Hubble rate H . Since $H \propto T^2/m_P$, with T the temperature, there is an absolute energy scale given by the Planck mass m_P : the lower the scale of baryogenesis compared to m_P , the harder it becomes to have at the same time a large enough CP asymmetry and small washouts.

In the IDM it was shown [10] that this intimate connection between CP asymmetry and washouts precludes leptogenesis for, roughly, $M \equiv M_1 \lesssim O(10^2)$ TeV, unless one of the low-scale-baryogenesis mechanisms described below is used. Arguably this bound is quite model independent [10] (within an order of magnitude, check e.g. [11] for the analogous result in the inverse seesaw²). The importance of this type of washouts also becomes apparent comparing the resultant baryon asymmetry after including them or not, see figure 2.

Next we briefly describe the three mechanisms which, in our opinion, have been demonstrated to allow for baryogenesis at the TeV scale -or even below-.

- *Almost degenerate particles:* When there is a pair of almost degenerate particles, the CP asymmetry in decays can be enhanced up to $O(1)$ values [13, 14]. To be more concrete consider two sterile neutrinos N_i ($i = 1, 2$), with masses M_i , Yukawa couplings $\lambda_{\alpha i}$ with the SM lepton doublet ℓ_α , and decay widths Γ_i . If the resonant condition $\Delta M \equiv M_2 - M_1 = \Gamma_2/2$ holds, then the CP asymmetry in N_1 decays, ϵ_1 , can take values as large as $1/2$ independently of the size of the Yukawa couplings of N_2 , i.e. of $(\lambda^\dagger \lambda)_{22}$. Moreover, for $\Gamma_{1,2} \ll \Delta M \ll M_1$, $\epsilon_1 \propto (\lambda^\dagger \lambda)_{22}/\delta$, with $\delta \equiv \Delta M/M_1$. Therefore it is possible to reduce the “dangerous” washouts

taking $(\lambda^\dagger \lambda)_{22}$ small enough, while keeping ϵ_1 sizable by choosing a tiny value for δ . This -so called resonant leptogenesis- mechanism has been widely studied (see e.g. [15]) and it can lead to successful baryogenesis at the TeV scale. The amount of degeneracy required has been studied in detail in [10] for the case of sterile neutrinos. It depends on M_1 and on the hierarchy among the Yukawa couplings, r , defined as the minimum non-null quantity in the list $\{\sqrt{(\lambda^\dagger \lambda)_{11}/(\lambda^\dagger \lambda)_{22}}, \sqrt{K_{\alpha i}} (i = 1, 2; \alpha = e, \mu, \tau)\}$, where the projectors are given by $K_{\alpha i} \equiv \frac{|\lambda_{\alpha i}|^2}{(\lambda^\dagger \lambda)_{ii}}$. E.g. the following rules of thumb apply to get the BAU via the resonant leptogenesis mechanism:

$$\delta \cdot r \lesssim 10^{-8} \text{ for } M_1 \sim 4 \text{ TeV},$$

$$\delta \cdot r \lesssim 2 \times 10^{-9} \text{ for } 250 \text{ GeV} \lesssim M_1 \lesssim 1 \text{ TeV}.$$

- *Late decay:* The rate of the washout processes associated to the CP asymmetry, like $H_2 \ell \leftrightarrow \bar{H}_2 \bar{\ell}$ in figure 1, is proportional to $(T/M_2)^a$ when $T \lesssim M_2$, with M_2 the mass of the mediator and $a = 2$ (4) for a Majorana (Dirac or scalar) propagator. Increasing the value of M_2 relative to M_1 (the mass of the decaying particle N_1) does not help, of course, to lower the scale of baryogenesis, because the CP asymmetry and washouts would be reduced by the same amount. However the later the N_1 ’s decay, the smaller the washouts will be at the time the baryon (or lepton) asymmetry is generated. Successful baryogenesis at the TeV scale via this late-decay scenario requires two conditions: a small decay width $\Gamma_1 \ll H(T = M_1)$ and a new process to create the N_1 ’s (given that the inverse decays and related scatterings are tiny), e.g. an exotic gauge interaction [16, 17]. Note that the “new” process must decouple before N_1 decays, so that the N_1 ’s disappear via the CP-violating interaction. In this way it is possible to have leptogenesis at $T \sim$ few TeVs (the exact value depending on the freeze-out temperature of the sphalerons) and baryogenesis at much lower scales if B is violated perturbatively.
- *Massive decay or annihilation products:* It has been explained that the CP asymmetry is proportional to the amplitude of processes that violate B (or L) and which therefore contribute to the washout of the baryon asymmetry. The rate of the washouts is not only determined by the amplitude of the corresponding process, but also by the number density of the particles involved in the interac-

¹To simplify the notation we will often omit flavour indices.

²A quite different scenario involving CP-violating decays and annihilations has been presented in [12]. Although the parameter space was not completely scanned, the results also suggest a bound $M \gtrsim O(10^2)$ TeV for the scale of baryogenesis (when none of the low-scale-baryogenesis mechanisms described below are implemented).




Figure 1: The CP asymmetry in N_1 decays comes -at leading order- from the interference of the tree-level and one-loop diagrams shown on the top. The vertical cuts through the loops indicate that those particles can go on shell, which is necessary to have a CP even phase. In turn, this implies the existence of the processes on the bottom, which contribute to the washout of the lepton asymmetry.




Figure 2: The final baryon asymmetry, Y_B , as a function of M , including the washout processes $H_2\ell \leftrightarrow \bar{H}_2\bar{\ell}$ and $\ell\ell \leftrightarrow \bar{\ell}\bar{H}_2$ (solid blue curve) or not including them (dashed red curve). The Yukawa couplings and M_2 are chosen so that $\frac{\Gamma_1}{H(T=M)}$ and ϵ keep constant for all M , where Γ_1 is the decay width of N_1 . As can be seen, for $M \lesssim 10^7$ GeV the baryon asymmetry falls exponentially with decreasing M . Instead, if the washouts are -incorrectly- not included, Y_B only declines linearly and its value for M at the TeV scale is wrong by many orders of magnitude.

tion. If at least one of these particles is massive and becomes non-relativistic during the baryogenesis epoch, the corresponding washout rate will be Boltzmann suppressed. This mechanism was shown to allow for TeV scale baryogenesis from DM annihilation first in [18] and from heavy particle decays in [10].

Actually the full mechanism is somewhat more complicated. To see why let us consider again the

IDM as an example, with H_2 playing the role of the massive decay product. The decays of N_1 generate an asymmetry not only in leptons, but also in H_2 . It is a simple statistical fact that the asymmetries in both sectors, ℓ and H_2 , induce washouts of the lepton asymmetry (e.g. more ℓ 's than $\bar{\ell}$'s, or more H_2 's than \bar{H}_2 's, favors the occurrence of $\ell H_2 \rightarrow \bar{\ell} \bar{H}_2$ over $\bar{\ell} \bar{H}_2 \rightarrow \ell H_2$). At linear order in the asymmetries $Y_{\Delta X} \equiv Y_X - Y_{\bar{X}} \equiv \frac{n_X}{s} - \frac{n_{\bar{X}}}{s}$ (and assuming kinetic equilibrium),

$$\frac{dY_{\Delta\ell}}{dz} = \frac{-1}{sHz} \left\{ \frac{Y_{\Delta\ell}}{Y_{\ell}^{\text{eq}}} + \frac{Y_{\Delta H_2}}{Y_{H_2}^{\text{eq}}} \right\} \gamma^{\text{eq}}(\ell H_2 \leftrightarrow \bar{\ell} \bar{H}_2) + \dots,$$

where $z \equiv M_1/T$ and the superscript “eq” indicates use of equilibrium phase space distributions with zero chemical potentials. To appreciate the washout effects on $Y_{\Delta\ell}$, write the i th term of the RHS of this equation as $-Y_{\Delta\ell} w_i(z)$ and analyze the function $w_i(z)$. The first term corresponds to a washout $\propto \gamma^{\text{eq}}(\ell H_2 \leftrightarrow \bar{\ell} \bar{H}_2)/Y_{\ell}^{\text{eq}}$, which is clearly Boltzmann suppressed by $e^{-M_{H_2}/T}$. The problem arises with the second one: how is $Y_{\Delta H_2}$ related to $Y_{\Delta\ell}$? If they were linked by a conservation law, so that $Y_{\Delta H_2} = \text{number} \times Y_{\Delta\ell}$, then $w_2(z) \propto \gamma^{\text{eq}}(\ell H_2 \leftrightarrow \bar{\ell} \bar{H}_2)/Y_{H_2}^{\text{eq}}$, which is NOT Boltzmann suppressed (as long as the mass of ℓ is negligible). In this case the mechanism does NOT work to lower the scale of baryogenesis. Instead if H_2 can disappear through some fast interaction different from the one at the origin of the BAU, the corresponding chemical equilibrium condition would imply that $Y_{\Delta H_2}$ is Boltzmann suppressed with respect to

$Y_{\Delta\ell}$. Then it is possible to achieve leptogenesis at the TeV scale (or, if B is violated perturbatively, baryogenesis at much lower temperatures).

The IDM is very useful to illustrate this issue. One of the allowed terms in the scalar potential is $\lambda_5(H^\dagger H_2)^2/2$. When the coupling $\lambda_5 = 0$ a conserved “lepton” number can be defined, assigning $L = -1$ to H_2 . Instead for large λ_5 there are fast “lepton” number violating interactions, such as $H_2\bar{H} \leftrightarrow \bar{H}_2H$. The amount of baryon asymmetry that can be generated at the TeV scale as a function of λ_5 is plotted in figure 3, which should serve as a lesson of when it is possible and when it is not possible to have baryogenesis at the TeV scale using massive decay or annihilation products. Later we will review some variations on this mechanism exemplified with models of baryogenesis from DM annihilation.




Figure 3: The final baryon asymmetry, Y_B , as a function of λ_5 in the IDM with $M_1 = 2.8$ TeV and $M_{H_2} = 0.6M_1$. The freeze-out temperature of the sphalerons is taken equal to $T_* \simeq 130$ GeV [19]. The Yukawa couplings and M_2 have a fix value which is not important for the discussion. It is clear that the BAU, $Y_B^{\text{obs}} \sim 10^{-10}$, can only be obtained for large values of λ_5 , so that the processes $H_2\bar{H} \leftrightarrow \bar{H}_2H$ are fast during the relevant period for baryogenesis.

We have seen that the IDM is a simple extension of the SM which allows to incorporate any of the three low-scale-baryogenesis mechanisms described above ³ [10] and it also provides a DM candidate, H_2 . A similar situation occurs when extending the SM with three sterile neutrinos, N_i , and a charged scalar, δ^+ , singlet under SU(2) [20] (in this case the DM is N_1 , with $M_1 > \text{few KeV}$ s and the BAU can be generated from the

CP-violating decays of N_2). To finish this very short review we wish to quote an example, for each of the three mechanisms, of their implementation in models aimed to explain the similar abundances of baryonic and dark matter (at least up to some degree) and that preserve the “WIMP miracle”.

In [21] baryogenesis occurs via leptogenesis in the decay of some gauge singlet Dirac fermions Ψ with $O(1)$ TeV masses. The density of Ψ when the baryon asymmetry starts to build up has been established by the freeze out of annihilations, so that the baryon density asymmetry is similar to the relic density of a Weakly Interacting Massive Particle (WIMP) if the CP asymmetry is $O(1)$. Then a pair of almost degenerate Ψ ’s are required to have a large CP asymmetry and leptogenesis at the TeV scale.

The late decay mechanism was incorporated in [22], where the scale of baryogenesis is only constrained to be above that of Big Bang nucleosynthesis. In this scenario there are at least two species of WIMPs, χ_{DM} which is stable DM, and χ_B which is meta-stable and decays after its annihilations freeze out. The decay violates CP and B , generating an amount ϵ of baryon asymmetry per decay. Since the χ_B ’s start to decay at $T \ll m_{\chi_B}$, baryogenesis is safe from the type of dangerous washouts depicted in figure 1. Moreover, the population of χ_B ’s has a WIMP-like relic density at the start of baryogenesis, which was established by the freeze out of interactions different from the CP-violating ones. Altogether, there is a simple relation among the relic abundances of baryons (Ω_B) and DM ($\Omega_{\text{DM}} = \Omega_{\chi_{\text{DM}}}$):

$$\frac{\Omega_B}{\Omega_{\text{DM}}} \simeq \epsilon \frac{m_p}{m_{\chi_B}} \frac{\tilde{\Omega}_{\chi_B}}{\Omega_{\chi_{\text{DM}}}},$$

where m_p is the proton mass, m_{χ_B} the χ_B mass, and $\tilde{\Omega}_{\chi_B}$ the would-be relic abundance of χ_B were it stable.

Finally let us consider baryogenesis from DM annihilations, which is a quite minimalistic way of relating the BAU and DM while preserving the WIMP miracle, because the baryon asymmetry is generated in the same annihilations whose freeze out determine the abundance of DM. Here a massive annihilation product Ψ is used to allow for baryogenesis at a low scale -set by the mass of the DM, m_χ -. Then the DM χ annihilates via $\chi\chi \rightarrow \Psi a, \bar{\Psi}\bar{a}$, where a is some light -or massless- field which accumulates particle-antiparticle asymmetry due to the CP-violating annihilations. This asymmetry might be baryonic or of some other type (like leptonic), in which case it must be transferred to the baryon sector by some fast process (like the electroweak sphalerons). In turn the asymmetry in the heavy Ψ ’s

³The realization of the late-decay scenario requires some extra way of producing the heavy neutrinos.

could induce fatal washouts, as explained before. To avoid this problem different ideas have been proposed.

In [18] Ψ is an exotic vector-like lepton or quark that decays due to fast interactions involving light sterile particles (see also [23]). These particles belong to a dark sector decoupled from the SM fields at low temperatures. Both models with and without a conserved $B-L$ were presented in [18] (in the former case the asymmetry in Ψ is transferred to the light sterile sector avoiding a complete cancellation of the baryon asymmetry after Ψ decays). Then in [24] it was shown that successful baryogenesis is possible without the light hidden sector, even in models with a conserved $B-L$. Here Ψ , a vector-like quark, decays fast into an exotic particle Ψ_2 , that in turn decays into SM particles after the electroweak sphalerons freeze out (hence the final baryon asymmetry is equal to the lepton asymmetry that keeps frozen after the sphalerons decouple).

A different approach to solve the problem associated with the asymmetry in the heavy annihilation (or decay) product was proposed in [25], namely to make a Majorana fermion or real scalar play the role of Ψ . In this way there is no asymmetry accumulating in the heavy field and the aforementioned problem is completely avoided. An implementation of this idea developed in [25] is to have baryogenesis from DM annihilation, like in the models just described, but the DM annihilates via $\chi\chi \rightarrow N\nu$, where N is a heavy Majorana neutrino with $M = O(1)$ TeV. The DM χ is a singlet fermion and therefore ν must also be a sterile neutrino (with mass m_ν). As long as $m_\nu \ll T$, an helicity asymmetry might be generated in the annihilations. The helicity asymmetry can be partially transferred to the SM lepton sector through fast Yukawa interactions and subsequently to baryons via the electroweak sphalerons. Therefore the BAU freezes in when the sphalerons decouple at $T_* \sim 100$ GeV and the condition for m_ν becomes $m_\nu \ll T_*$. In this model the sterile neutrinos are responsible for the masses of the SM neutrinos and they play a key role in baryogenesis and the freeze out of DM annihilations.

Acknowledgments

I wish to thank my collaborators on this subject, Nicolás Bernal, Stefano Colucci, François-Xavier Josse-Michaux, Nuria Rius, and Lorenzo Ubaldi.

This work has been supported by the Spanish MINECO Subprogramme Juan de la Cierva and it has also been partially supported by the Spanish MINECO grants FPA2011-29678-C02-01 and Consolider-Ingenio CUP (CSD2008-00037), and by Generalitat Valenciana

grant PROMETEO/2009/116. In addition we acknowledge partial support from the European Union FP7 ITN INVISIBLES (Marie Curie Actions, PITN- GA-2011-289442).

References

- [1] M. Y. Khlopov and A. D. Linde, *Is It Easy to Save the Gravitino?*, *Phys. Lett.* **B138** (1984) 265–268.
- [2] J. R. Ellis, J. E. Kim, and D. V. Nanopoulos, *Cosmological Gravitino Regeneration and Decay*, *Phys. Lett.* **B145** (1984) 181.
- [3] F. Vissani, *Do experiments suggest a hierarchy problem?*, *Phys. Rev.* **D57** (1998) 7027–7030, [[hep-ph/9709409](#)].
- [4] J.-M. Frère, T. Hambye, and G. Vertongen, *Is leptogenesis falsifiable at LHC?*, *JHEP* **0901** (2009) 051, [[arXiv:0806.0841](#)].
- [5] F. F. Deppisch, J. Harz, and M. Hirsch, *Falsifying High-Scale Leptogenesis at the LHC*, *Phys. Rev. Lett.* **112** (2014) 221601, [[arXiv:1312.4447](#)].
- [6] S. Davidson and A. Ibarra, *A lower bound on the right-handed neutrino mass from leptogenesis*, *Phys. Lett.* **B535** (2002) 25, [[hep-ph/0202239](#)].
- [7] T. Hambye, Y. Lin, A. Notari, M. Papucci, and A. Strumia, *Constraints on neutrino masses from leptogenesis models*, *Nucl. Phys.* **B695** (2004) 169, [[hep-ph/0312203](#)].
- [8] D. V. Nanopoulos and S. Weinberg, *Mechanisms for Cosmological Baryon Production*, *Phys. Rev.* **D20** (1979) 2484.
- [9] E. Ma, *Verifiable radiative seesaw mechanism of neutrino mass and dark matter*, *Phys. Rev.* **D73** (2006) 077301, [[hep-ph/0601225](#)].
- [10] J. Racker, *Mass Bounds for Baryogenesis from Particle Decays and the Inert Doublet Model*, *JCAP* **1403** (2014) 025, [[arXiv:1308.1840](#)].
- [11] J. Racker, M. Peña, and N. Rius, *Leptogenesis with small violation of B-L*, *JCAP* **1207** (2012) 030, [[arXiv:1205.1948](#)].
- [12] I. Baldes, N. F. Bell, A. Millar, K. Petraki, and R. R. Volkas, *The role of CP violating annihilations in baryogenesis - case study of the neutron portal*, [arXiv:1410.0108](#).
- [13] M. Flanz, E. A. Paschos, U. Sarkar, and J. Weiss, *Baryogenesis through mixing of heavy Majorana neutrinos*, *Phys. Lett.* **B389** (1996) 693–699, [[hep-ph/9607310](#)].
- [14] L. Covi and E. Roulet, *Baryogenesis from mixed particle decays*, *Phys. Lett.* **B399** (1997) 113–118, [[hep-ph/9611425](#)].
- [15] A. Pilaftsis and T. E. Underwood, *Resonant leptogenesis*, *Nucl. Phys.* **B692** (2004) 303–345, [[hep-ph/0309342](#)].
- [16] M. Plümacher, *Baryogenesis and lepton number violation*, *Z. Phys.* **C74** (1997) 549, [[hep-ph/9604229](#)].
- [17] J. Racker and E. Roulet, *Leptogenesis, Z-prime bosons, and the reheating temperature of the Universe*, *JHEP* **0903** (2009) 065, [[hep-ph/0812.4285](#)].
- [18] Y. Cui, L. Randall, and B. Shuve, *A WIMPy Baryogenesis Miracle*, *JHEP* **1204** (2012) 075, [[arXiv:1112.2704](#)].
- [19] M. D’Onofrio, K. Rummukainen, and A. Tranberg, *The Sphaleron Rate in the Minimal Standard Model*, *Phys. Rev. Lett.* **113** (2014) 141602, [[arXiv:1404.3565](#)].
- [20] M. Frigerio and C. E. Yaguna, *Sterile Neutrino Dark Matter and Low Scale Leptogenesis from a Charged Scalar*, [arXiv:1409.0659](#).
- [21] S. Davidson and M. Elmer, *Similar Dark Matter and Baryon abundances with TeV-scale Leptogenesis*, *JHEP* **1210** (2012) 148, [[arXiv:1208.0551](#)].

- [22] Y. Cui and R. Sundrum, *Baryogenesis for weakly interacting massive particles*, *Phys.Rev.* **D87** (2013) 116013, [[arXiv:1212.2973](#)].
- [23] N. Bernal, F.-X. Josse-Michaux, and L. Ubaldi, *Phenomenology of WIMPy baryogenesis models*, *JCAP* **1301** (2013) 034, [[arXiv:1210.0094](#)].
- [24] N. Bernal, S. Colucci, F.-X. Josse-Michaux, J. Racker, and L. Ubaldi, *On baryogenesis from dark matter annihilation*, *JCAP* **1310** (2013) 035, [[arXiv:1307.6878](#)].
- [25] J. Racker and N. Rius, *Helicitogenesis: WIMPy baryogenesis with sterile neutrinos and other realizations*, [arXiv:1406.6105](#).